Predictors of mental health and behavioural crises in children and youth with neurodevelopmental disorders

Lenz, S. [1], Rapley, J. [1], Canning, E. [1, 2], Kushki, A. [1], Baribeau, D. [1] **1) Bloorview Research Institute** 2) Carleton University

Higher rates of crises,

resulting in hospital

visits & police calls²

Background

Children and youth with neurodevelopmental disorders (NDDs) experience:




Increased incidence of aggression, self-injury, & suicidal ideation

Research Question

What are the predictors of mental health/behavioural crises in children and youth with NDDs?


Methods




Designed survey to collect patient demographic, clinical, and environmental data

Collected longitudinal data from 270 psychopharmacology clinic patient charts





Performed cox proportional hazards regressions to identify predictors of mental health and behavioural crises



1. Giannouchos TV, Beverly J, Christodoulou I, Callaghan T. Suicide and non-fatal self-injury-related emergency department visits among individuals with autism spectrum disorder. Autism 2023;27(7):1983-1996. DOI:10.1177/13623613221150089 2.Schott W, Tao S, Shea L. Emergency Visits for Autistic Children and Children With ADHD. Pediatrics 2022;149(Suppl 4):e2020049437V. DOI:10.1542/peds.2020-049437V

Social factors significantly predict mental health & behavioural crises in children and youth with neurodevelopmental disorders





Holland Blcorview

Kids Rehabilitation Hospital Foundation